



The Hub Dental Practice



Love Your Smile

775 Witan Gate, Central Milton Keynes MK9 2FW

Love Your Smile

01908 690326

reception@TheHubDentalPractice.com

www.TheHubDentalPractice.com

Dr David Gilmartin,
 Dr Kamilah Makda
 Dr Chivani Tailor,
 Dr Ivo Zdravkov,
 Dr Allen Betsis,
 Dr Etienne Deysel, Sedationist

Dr Aaron Lopez, Specialist Periodontist
 Dr Monica Cueva Moya, SI Periodontics
 Dr Gulelala Azhar, SI Restorative,
 Dr Giovanni Marras, PLT Periodontics
 Miss Sam Singleton, Practice Manager
 Mrs. Nur Gilmartin, Dental Hygienist

Dr Alvaro De Castro Torres, SI Dental Implants & RCT
 Dr Konstantinos Tzamalass, Specialist Orthodontist
 Dr Iyad Abou-Rabii, PLT Dental Implants & Oral Surgery
 Dr Ezgi Cilingir, PLT Dental Implants & Periodontics
 Dr Jonathan Harper, PLT Endodontics
 Dr Jaime De Castro Torres, SI Dental Implants

Apicectomy


The Problem

An infection has occurred at the tip of the root or roots of a tooth, spreading into the bone that supports the tooth. Occasionally people may be unaware of the problem, but usually there is discomfort, episodes of swelling, gumboils or bad taste.

Why do I need Treatment?

If left untreated the infection will spread, possibly developing into an abscess or cyst, or damaging bone around adjacent teeth. The tooth may become loose.

The infection cannot be cured with antibiotics, but they are often used to give temporary relief of symptoms.



What is the Treatment?

An Apicectomy is a small surgical procedure designed to cure an infection at the tip of the root or roots of a tooth. It is indicated when, for a variety of reasons, it has proved impossible for your dentist to cure the infection by removing the dead nerve and placing a root filling. It is most frequently carried out under local anaesthetic (injection into the gum).

A small cut is made in the gum, which is then lifted off the bone. Access through the bone to the root tip is made using a drill.

The infection is cleaned out, part of the root tip removed and a small filling placed in the end of the root. The gum is replaced with dissolvable stitches.

What can I expect after the operation?

As the local anesthetic wears off a few hours after surgery there will be some discomfort, but this is quite variable. This is usually managed with an analgesic such as Ibuprofen. Antibiotics may also be prescribed. Some swelling at the operation site and of the overlying skin is normal.

This is worse 24-48 Hours after the operation and resolves over the next few days. It is important to keep the site of surgery as clean as possible.

The area can be brushed gently with a toothbrush softened in hot water, together with hot salty mouthwash (teaspoon of salt in a beaker of warm water), 2-3 times a day, beginning the day after surgery and continuing for 1 week.

What are the possible problems?

Prolonged bleeding is rare, but if it occurs then pressure with a cotton handkerchief or swab for at least 10 minutes usually stops it.

Occasionally the gum will shrink back slightly following surgery, which may expose the margin of a crown if one is present.

The operation is occasionally unsuccessful at removing the infection. If unsuccessful after 2 attempts the tooth is probably better removed.

Some lower tooth roots lie very close to important nerves which if damaged may cause numbness of the lower lip and chin on one side.

If this risk is present your surgeon will discuss it with you.

Dr Iyad Abou-Rabii



Dr Iyad Abou-Rabii offers a full range of Oral Surgery and Implant treatments.

Iyad is an extremely experienced surgeon with an extensive experience in general dental surgery and Dental Implants.

Iyad has completed more than 2000 implants, and 1000's of minor oral surgery procedures (impacted teeth).

Iyad has published many articles in international Dental and Medical Journals