



# The Hub Dental Practice



Love Your Smile

775 Witan Gate, Central Milton Keynes MK9 2FW

Love Your Smile

01908 690326

[reception@TheHubDentalPractice.com](mailto:reception@TheHubDentalPractice.com)

[www.TheHubDentalPractice.com](http://www.TheHubDentalPractice.com)

Dr David Gilmartin,  
Dr Kamilah Makda  
Dr Chivani Tailor,  
Dr Ivo Zdravkov,  
Dr Allen Betsis,  
Dr Etienne Deysel, Sedationist

Dr Aaron Lopez, Specialist Periodontist  
Dr Monica Cueva Moya, SI Periodontics  
Dr Gulelala Azhar, SI Restorative,  
Dr Giovanni Marras, PLT Periodontics  
Miss Sam Singleton, Practice Manager  
Mrs. Nur Gilmartin, Dental Hygienist

Dr Alvaro De Castro Torres, SI Dental Implants & RCT  
Dr Konstantinos Tzamalass, Specialist Orthodontist  
Dr Iyad Abou-Rabii, PLT Dental Implants & Oral Surgery  
Dr Ezgi Cilingir, PLT Dental Implants & Periodontics  
Dr Jonathan Harper, PLT Endodontics  
Dr Jaime De Castro Torres, SI Dental Implants

## Peri-Implantitis Surgery Information and Consent

Periodontal surgery may be seen as a secondary line of treatment of those pockets persisting after initial treatment. These are often the area's most severely affected by ... periodontal disease. Surgery has the advantage of allowing direct access, inspection and cleaning of the affected root surfaces. Many long term periodontal studies have demonstrated the benefit and success of periodontal surgery.

### Benefits of Peri-Implantitis Surgery

- More predictable results in reducing pocketing and inflammation
- Reduce the possibility of future bone support loss, and therefore tooth loss
- Can help to allow for easier cleaning
- Able to investigate the areas with inflammation and pocketing, and remove diseased tissue

### Aims of Peri-Implantitis Surgery

Periodontal surgery is primarily carried out to eliminate/reduce pockets, and reduce bacterial load. The plaque growing on the implant surfaces is the causative factor for periodontal disease, and associated bone support loss. Clean implant surfaces leads to healthy gum tissue.

It is also carried out to provide a more favourable long term outcome.

### Peri-Implantitis Surgery

After the area is completely numbed an incision is made to allow the gum tissue to be lifted away from the implant.

The inflamed areas of the inflamed gum are removed and the implant surface is thoroughly cleaned. The bone may need to be resected. The gum tissue may need to be recontoured to better adapt to the neck of the tooth. The gum is then replaced, usually involving stitches and a dressing.

### Alternative to suggested treatment

I understand that alternatives to periodontal surgery include: no treatment-with the expectation of possible advancement of my condition which may result in permanent loss of teeth: extraction of teeth involved with periodontal disease; and non-surgical scraping of tooth roots and lining of the gum (scaling and root planning), with or without medication, in an attempt further, to reduce bacteria and tartar under the gum-line

However this may not fully eliminate deep bacteria and tartar, may not reduce gum pockets, will require more frequent professional care and time commitment and may result in worsening of my condition and the premature loss of teeth.

In some cases medications and/or other materials and/or tissue grafts may be used during the procedure. In this case a separate Consent form will be used.

### Side Effects

Following on from periodontal surgery there can be swelling, bruising and discomfort around the treated area. This usually settles within 3-5 days.

There can be more dental hypersensitivity (this means that you can have pronounced sensation to hot and cold). This sensitivity is usually short-term, and can be relieved by desensitising tooth pastes.

The teeth being operated on can appear longer. This can be aesthetically significant if surgery is being performed on teeth at the front of the mouth.

Occasionally the teeth may become more mobile and suffer of aesthetic exposure of margins.

Infections and necrosis of the tissues may occur. In this situation the use of antibiotics is required to reduce the load of bacteria. Nevertheless, a necrotic site requires several months to be totally healed.

Profuse bleeding is not so common and is normally treated with haemostatic agents in the dental environment.

## Success

The long term success of any periodontal procedure depends on a variety of factors. This includes the extent of disease on initial presentation, the complexity of the lesions that are being operated on.

Effective plaque control is vital, and you will need to continue to maintain a good standard of oral hygiene.

Avoiding smoking is also very important especially in the early days of healing. Smokers have delayed healing and a less favourable response to treatment.

Generalised risk factors are smoking, and uncontrolled diabetes.

Other very important factors are maintaining a high standard of oral hygiene, and regular maintenance.

## LONG-TERM CARE

Many authorities have demonstrated that a long term maintenance programme is the most important predictor of success in periodontal disease treatment outcome. The most important factor in maintaining gum health is the daily removal of plaque from the tooth-gum junction by the patient. Most patients find it necessary to see a dentist/hygienist at regular intervals to ensure that a good standard of oral hygiene is maintained and to monitor gum health, enabling any problems to be identified and dealt with early. We recommend periodontal maintenance on a three monthly basis for the first year after surgery.

## Informed consent

**PATIENT CONSENT: by my signature below, I expressly acknowledge that:**

The dentist has explained the likely outcomes and possible complications of each alternative option.

The dentist has made leaflets and other relevant information available to me to help with my decision making.

The dentist has given me an opportunity to ask questions about any matters related to my treatment, raise any other concerns and given me the opportunity to postpone the treatment or seek a second opinion.

The dentist has explained any significant risks and problems associated with Gum Surgery specific to me, and the likely outcomes if complications occur.

The dentist has explained the options for anaesthesia (local, sedation, general) and problems specific to each choice, and the likely outcomes if complications occur.

**The Dentist has explained to me that the procedure can be stopped at any time. The patient only needs to say stop, put up a hand, shout or make another signal etc. We say again, the patient can decide to stop the procedure at any time and for whatever reason they wish. We will always respect the patient's decision.**

**The dentist has explained the importance of following the post Treatment instructions and any additional instructions that are given verbally.**

The dentist has explained the options for waiting before I make my decision.

You have given us an up to date account of your medical and dental history, especially about any recent changes.

**You understand the risks of the procedure, including the risks that are specific to me, and the likely outcomes. The dentist has explained other relevant treatment options and their associated risks. The dentist has explained my prognosis and the risks of not having the procedure.**

You were able to ask questions and raise concerns with the dentist about my condition, the procedure and its risks, and my treatment options. And that your questions and concerns have been discussed and answered to your complete satisfaction.

You understand that other dental procedures may be done if further dental disease is found during the procedure, or to correct other problems in my mouth. And that you understand that if teeth are removed during the dental treatment, that these may be retained for training purposes and then disposed of sensitively.

You understand that no guarantee has been made that the procedure will improve the condition, and may even make your condition worse.

**On the basis of the above statements, I request to have the Peri-Implantitis Surgery described above.**

**I realise that signing this does not oblige me to have all or any part of the treatment proposed, but is simply an acknowledgement of the fact that I am giving my informed consent if I decide to go ahead.**

**(Peri-Implantitis Surgery)**

Name of Patient: \_\_\_\_\_ Date of Birth: \_\_\_\_//\_\_\_\_//\_\_\_\_

Signature: \_\_\_\_\_ Date: \_\_\_\_//\_\_\_\_//\_\_\_\_