



Love Your Smile

The Hub Dental Practice

775 Witan Gate, Central Milton Keynes MK9 2FW



Love Your Smile

01908 690326

reception@TheHubDentalPractice.com

www.TheHubDentalPractice.com

Dr David Gilmartin,
Dr Kamilah Makda
Dr Chivani Tailor,
Dr Ivo Zdravkov,
Dr Allen Betsis,
Dr Etienne Deysel, Sedationist

Dr Aaron Lopez, Specialist Periodontist
Dr Monica Cueva Moya, SI Periodontics
Dr Gulelala Azhar, SI Restorative,
Dr Giovanni Marras, PLT Periodontics
Miss Sam Singleton, Practice Manager
Mrs. Nur Gilmartin, Dental Hygienist

Dr Alvaro De Castro Torres, SI Dental Implants & RCT
Dr Konstantinos Tzamalass, Specialist Orthodontist
Dr Iyad Abou-Rabii, PLT Dental Implants & Oral Surgery
Dr Ezgi Cilingir, PLT Dental Implants & Periodontics
Dr Jonathan Harper, PLT Endodontics
Dr Jaime De Castro Torres, SI Dental Implants

Removal of Wisdom Teeth

This information is for patients who may need to have an operation to remove their impacted wisdom teeth. It explains why they may need to be removed, what is involved and any risks or complications that there may be.

The wisdom teeth

Adults normally have 32 teeth, and wisdom teeth (3rd molars) are the last to come through at the back of the mouth. Normally there are 4 wisdom teeth, one in each side of the upper and lower jaw.

Impacted wisdom teeth

If the jaws are too small to accommodate all the teeth, there may not be enough space for the wisdom teeth to come through properly and they become impacted (stuck), often causing problems.

Reasons for the removal of wisdom teeth

1. The most common reason is recurrent infection of the gum overlying a tooth that is part way through the gum (pericoronitis).
2. Decay (caries) in the wisdom tooth, which your dentist cannot restore.
3. Infection of the tooth (abscess) due to advanced dental decay.
4. When the adjacent molar tooth is affected by gum (periodontal) disease or dental decay due to the impacted wisdom tooth.
5. Progressive cystic (fluid filled sac) formation around the tooth.
6. As part of other surgical procedures involving the jaw.
7. There may be other less common reasons that your surgeon will discuss with you.

The removal of wisdom teeth

- There is great variation in the difficulty of removing wisdom teeth.
- The procedure can be carried out under local anaesthesia (injection in the gum to numb the area), with or without intravenous sedation (injection in the arm or hand to reduce anxiety), or under general anaesthetic (completely asleep in a hospital).

- Your surgeon will discuss with you which method is most appropriate.
- The degree of difficulty of the surgery, any underlying medical conditions and other personal circumstances will be taken into account choosing the method.
- The procedure can involve an incision (cut) in the gum close to the tooth.

Sometimes some jaw bone around the tooth is removed with a drill and the procedure can often be made easier by sectioning (cutting up) the tooth itself into smaller pieces.

What can be expected after the operation?

- There is great variation in the pain and discomfort suffered by individuals and this also depends on how difficult the surgery is. The average recovery time is between 5-7 days.
- Swelling and bruising is common and tends to be at its worst on the 2nd and 3rd day after surgery reducing over about a week.
- Mouth opening is likely to be restricted and a semi-solid diet will be required for the first few days.
- There will be some pain from the operation site. Your surgeon will recommend the best pain killer to help relieve this after your operation.

Some possible complications

Removal of wisdom teeth is a very safe and common operation but there is some risk associated with most procedures:

- Most bleeding will have stopped shortly (within ½ hour) of the operation finishing. Blood stained saliva may be noticed for a day or two. More persistent bleeding may occur which should be dealt with by your surgeon/hospital.
- Sometimes antibiotics are prescribed after the operation. Even if they have infection of a tooth socket occasionally occurs.
- When an adjacent tooth has a large filling or crown it is possible that this can be dislodged during surgery.
- The nerves which give sensation to the lower lip and chin and to the tongue are very close to the lower wisdom teeth and can be prone to bruising.
- If this happens numbness or a tingling sensation in the lip and chin, or tongue will be experienced.
- This is temporary in most cases, but in a small number recovery may not be complete. Very rarely continued pain may be felt from damaged nerves.
- Your surgeon will inform you of other complications, which are relevant to your case.
- **We give you a consent form to read and sign, which explains the problems associated with Wisdom Tooth Extraction. It is essential to understand before you sign and we encourage questioning from patients.**
- **Both the consent form and Post Extraction leaflet are available on our website**

Please speak to your surgeon before your operation if you have any concerns about these risks.